



Apeldoornseweg 47 6814 BJ Arnhem
The Netherlands tel.: 0031 26 4427706
KvK-Arnhem nr 09099868 BTW: NL807257084B01

I-Health Protocols:

| Name: | Description: |
|---------------------------------|--|
| 1. CVA | <ul style="list-style-type: none">- Do an EDA to see where the energy gap is greatest;- Do an AFT-scan and press 'Get' in the analysis screen;- Use the chosen AFT-frequency according to the Mussat treatment schedule (use info-button in beamer and lightpen-screen);- underneath the feet (begin with 30 seconds during the first time) and at the back of the head (1 minute, gradually increasing).- Use the AFT-frequency at the affected limb or at the face for 5 or more minutes;- 2x / week.- Use cicatron with beamer at the region of the CVA. |
| 2. Benign hypertrophic Prostate | <ul style="list-style-type: none">- Have the patient sit on the beamer with the beamer as near to the prostate as possible;- The beamer is connected to the soundcard. The volume is set at 100% (as usual);- Use Emo-Chacratron 1 5- 10 minutes;- Use Prostanotron 5 – 10 minutes;- Use Prostatron 5- 10 minutes;- 2x tot 3x /week, total # 10 has shown very good results;- No measurements required other than 1 time at the beginning and one at the end for documentation. |
| 3. Lower Back Pain | <ul style="list-style-type: none">- EDA;- Vertebra Scan;- Use the combination of three signals, i-light when |



Apeldoornseweg 47 6814 BJ Arnhem
 The Netherlands tel.: 0031 26 4427706
 KvK-Arnhem nr 09099868 BTW: NL807257084B01

| | |
|-------------------------------------|---|
| | <p>available, as well as beamer 'Vertebra-signal', The related Emochacratron (1 or 2), Renastimutron;</p> <ul style="list-style-type: none"> - Primarily at the vertebra 50% of the treatment time, and secondly on the area of the dermatome when other symptoms require this. |
| <p>4. Claudicatio and open legs</p> | <ul style="list-style-type: none"> - EDA (will show lack of energy at lower meridians); - Do AFT-scan and select the appropriate frequency by pressing 'get'; - Treat the lower legs with the AFT signal (beamer attached to the i-health device); - Check that the liver and kidney can cope with the detoxification. May be it is necessary to add Carrot juice or another Vit A for the liver and a herbal tea for the kidney (or renastimutron directly on the kidneys); - 1x to t2x / week; - Check with Doppler. |
| <p>5. ME, CFS, Fibromyalgia</p> | <ul style="list-style-type: none"> - Eda; - Zapsan and use these frequencies for 2 weeks, then do another zapsan and use this. Zap 2x / week. - Check and correct acidity when necessary with baking soda, Megamin, barley grass or the like; - Scans: Sanoscan, weakest meridian scan, Vertebra-scan and treat according to the results with the beamer and the i-light. - The patient should drink enough water. Energize the water with detoxitron and let them take it home. |
| <p>6. Scars</p> | <ul style="list-style-type: none"> - Make an inventory of the scars; - Do an eda. - Use posttraumatron, emotraumatron when necessary, and lastly but very important: cicatron at the scars. |



Apeldoornseweg 47 6814 BJ Arnhem
The Netherlands tel.: 0031 26 4427706
KvK-Arnhem nr 09099868 BTW: NL807257084B01

| | |
|--|---|
| | <p>Lightpen as well as with the beamer.</p> <ul style="list-style-type: none">- Check with EDA. Document changes of scars. |
| 7. Allergies | <ul style="list-style-type: none">- Use the separate protocol (upon request participation in pilot). |
| 8. Food intolerances | <ul style="list-style-type: none">- Do EDA, provocation and another EDA;- Give the substance in the patient's hand and do another EDA;- Give the next substance, etc.;- I often ask the patient just to bring what is in their cupboard and eat regularly.- Evaluate by looking at the graphs and the test-results meta-analysis. |
| 9. Herpes infections, flu, and other systemic or intestinal infections | <ul style="list-style-type: none">- Use the zapper. |
| 10. Acne (bacterial) | <ul style="list-style-type: none">- When the lung meridian or lung-meridian and large intestine meridian are out of balance or blocked then use the zapper. (Otherwise it is also hormonal of origin and that is more complicated). |